



## Talk to your doctor about TAVNEOS® for your severe active GPA or MPA

### For about half of patients, even in remission, vasculitis symptoms are debilitating\*

\*According to an online, self-administered survey of 100 patients with GPA or MPA from July 21-August 25, 2022.

Having open, detailed conversations can help you and your doctor understand how severe active ANCA-associated vasculitis (GPA or MPA) is affecting you. Work together and decide if TAVNEOS® should be added to your treatment. Download, email, or take a screenshot of your guide and have it handy for your next appointment.

ANCA = anti-neutrophil cytoplasmic autoantibody; GPA = granulomatosis with polyangiitis, previously called Wegener's granulomatosis; MPA = microscopic polyangiitis.

#### 1 What symptoms have you been experiencing? Mark all that apply:

- Ears:** Hearing loss, ear infections, fluid leaking
- Eyes:** Irritated eyes, blurred vision
- Nose:** Crusted blood around nostrils, stuffy nose, sinus infection
- Lungs:** Trouble breathing, wheezing or cough
- Heart:** Chest pain
- Kidneys:** Blood in urine
- Digestive system:** Bloody diarrhea, stomach pain
- Skin:** Rashes, sore spots, blisters
- Nervous System:** Headache, numbness or tingling in the arms and legs
- General:** Muscle pain, joint pain

When did you first notice these symptoms? Have you had them before? Give examples and describe how you feel.

---

---

Be sure to tell your doctor about these and other symptoms you're having to help guide your care.

---

---

#### 2 Over the past few weeks, have your symptoms made it hard to work or do other activities? If so, describe here.

---

---

#### 3 How satisfied are you with your current treatment plan?



This guide is for anyone who has been diagnosed with severe active GPA or MPA, their loved ones, or those who are unsure if they have this disease.

Please see additional Important Safety Information on the next page and click here for the [Full Prescribing Information](#) and [Medication Guide](#).

#### WHAT IS TAVNEOS?

TAVNEOS is a prescription medicine that is used with other medicines (such as glucocorticoids) to treat adults with severe active anti-neutrophil cytoplasmic autoantibody (ANCA)-associated vasculitis (granulomatosis with polyangiitis [GPA], formerly known as Wegener's granulomatosis, and microscopic polyangiitis [MPA]).

**It is not known if TAVNEOS is safe and effective in children under the age of 18.**

#### IMPORTANT SAFETY INFORMATION

##### What is the most important information I should know about TAVNEOS?

TAVNEOS can cause serious side effects, including:

→ **Liver problems.** People taking TAVNEOS may have serious liver problems. Call your healthcare provider right away if you have unexplained symptoms such as:

- yellowing of your skin or the white part of your eyes (jaundice)
- pain on the upper right side of your stomach area (abdomen)
- feeling tired
- dark or brown (tea colored) urine
- bleeding or bruising more easily than normal
- loss of appetite

Your healthcare provider will do blood tests to check how well your liver is working before starting and during your treatment with TAVNEOS.

# Now that you've talked about how you're feeling, here are some questions to ask your doctor about TAVNEOS®:

- Do you think TAVNEOS® would be a good fit for me?
- How do I take TAVNEOS®?
- What are some common side effects of TAVNEOS®?

If TAVNEOS® is right for you, your doctor can fill out this [Start Form](#)

## IMPORTANT SAFETY INFORMATION (CONT'D)

### Do not take TAVNEOS if you are allergic to avacopan or any of the other ingredients in TAVNEOS.

→ Get medical help right away if you experience swollen lips, tongue, throat, trouble swallowing, or difficulty breathing. These could be signs of an allergic reaction. Do not take more TAVNEOS until you have consulted with your healthcare provider.

### Before taking TAVNEOS, tell your healthcare provider about all your medical conditions, including if you:

- have or have had abnormal liver blood tests.
- have or have had liver problems.
- have or think you may have hepatitis B or C.
- have an infection.
- are pregnant or are planning to become pregnant. It is not known if TAVNEOS will harm your unborn baby. Talk to your healthcare provider if you are pregnant or plan to become pregnant.
- are breastfeeding or plan to breastfeed. It is not known if TAVNEOS can pass into your breast milk. Talk to your healthcare provider about the best way to feed your baby if you take TAVNEOS.

**Tell your healthcare provider about all the other medicines you take,** including prescription and over-the-counter medicines, vitamins, and herbal supplements. TAVNEOS and certain other medicines may affect each other and cause side effects. Keep a list of the medicines you take and show it to your healthcare provider and pharmacist.

- Some medicines should not be taken with TAVNEOS.
- Your healthcare provider may prescribe other medicines to treat your disease.

### TAVNEOS may cause serious side effects, including:

- Liver problems (see above).

→ **Serious allergic reactions.** Stop taking TAVNEOS and get emergency medical help right away if you have any of the following signs of a serious allergic reaction: shortness of breath or trouble breathing; swollen lips, tongue, throat, or face; trouble swallowing; chest pain; feeling dizzy or faint; moderate or severe abdominal pain, or vomiting.

→ **Hepatitis B virus (HBV) reactivation.** If you have had HBV or are a carrier of HBV, taking TAVNEOS could cause the virus to become an active infection again. Tell your healthcare provider right away if you get worsening tiredness or yellowing of your skin or the white part of your eyes during treatment with TAVNEOS.

→ **Serious infections.** Serious infections can happen in people taking TAVNEOS, and these infections can lead to death. The most common serious infections with TAVNEOS were pneumonia and urinary tract infections. People with serious infections should not take TAVNEOS. Tell your healthcare provider right away if you have any symptoms of infection: fever, cold symptoms that do not go away, flu symptoms, pain during urination, or other signs of infection.

The most common side effects of TAVNEOS include:

- nausea, headache, high blood pressure, diarrhea, vomiting, rash, tiredness, stomach pain, dizziness, increase in blood creatinine, and burning or prickling sensation.

These are not all the possible side effects of TAVNEOS. Call your doctor for medical advice about side effects. Please see the [Full Prescribing Information](#) and [Medication Guide](#) for further details.

TAVNEOS is available as a 10 mg capsule.

You are encouraged to report negative side effects to Amgen at 1-833-828-6367 or to the FDA by visiting [www.fda.gov/medwatch](http://www.fda.gov/medwatch) or calling 1-800-332-1088.



© 2023 Amgen Inc. All rights reserved.  
08/23 USA-569-80326

Click here for the [Full Prescribing Information](#) and [Medication Guide](#).